

Rheumatoid arthritis with aquaporin-4 antibody-positive neuromyelitis optica receiving rituximab therapy

Chrong Reen Wang¹, Sheng-Min Hsu², Ying-Chen Chen³

¹Department of Internal Medicine, National Cheng Kung University Hospital, Tainan, Taiwan

²Department of Ophthalmology, National Cheng Kung University Hospital, Tainan, Taiwan

³Department of Medical Imaging, National Cheng Kung University Hospital, Tainan, Taiwan

A spectrum of autoimmune diseases has been associated with optic neuritis or myelitis in the presence of aquaporin-4 (AQP4) antibody, creating a classification named neuromyelitis optica spectrum disorder (NMOSD).¹ Further studies have identified a pathogenic role of B cells,² and rituximab (RTX) is an effective and safe treatment in this disease.³ Herein, we report a case of rheumatoid arthritis (RA) with anti-AQP4-positive optic neuritis. The RTX therapy resulted in absent autoantibody and clinical improvement.

A 58-year-old Han Chinese woman visited our hospital with a five-year history of polyarthritis and visual impairment. She was diagnosed to have RA with recent therapy including methotrexate (MTX) 10 mg/week and prednisolone 5 mg/day. For ocular presentations, there was retro-orbital pain with bilateral papillitis, followed by blurred vision with delayed responses/low amplitudes in visual evoked potential, leading to a diagnosis of optic neuritis. Nevertheless, she did not attend to the ophthalmological follow-up without taking medications, resulting in a progressive

eyesight loss to a counting fingers vision (10/5 cm, right/left).

Physical examination at the rheumatology clinic showed swollen, tender joints over wrist, metacarpophalangeal, proximal interphalangeal, and knee areas. Conventional X-ray demonstrated erosion over the wrist. Erythrocyte sedimentation rates (ESR, 52 mm/h) and C-reactive protein (CRP, 11.9 mg/L) levels were elevated with a 5.88 Disease Activity Score-28 (DAS28). Autoantibody profiles had increased anti-AQP4 (11 U/mL, enzyme-linked immunosorbent assay, RSR Limited, UK), rheumatoid factor (RF, 105 IU/mL) and anti-cyclic citrullinated protein (anti-CCP, 28 U/mL) levels. Evaluation for systemic lupus erythematosus (SLE) showed normal hemogram and negative urine analysis, as well as anti-nuclear antibody negativity, anti-double-stranded deoxyribonucleic acid, and anti-Sm antibodies. Sjögren's syndrome survey revealed negative SSA/SSB antibodies and minor salivary gland biopsy. There was no demyelination on brain/spinal cord magnetic

Received: January 05, 2021 **Accepted:** February 05, 2021 **Published online:** June 24, 2021

Correspondence: Chrong Reen Wang, MD, PhD. Department of Internal Medicine, National Cheng Kung University Hospital, 70403 Tainan, Taiwan.
Tel: 886-6-2353535 e-mail: z9109019@email.ncku.edu.tw

Citation:

Wang CR, Hsu SM, Chen YC. Rheumatoid arthritis with aquaporin-4 antibody-positive neuromyelitis optica receiving rituximab therapy. Arch Rheumatol 2022;37(1):153-155.

©2022 Turkish League Against Rheumatism. All rights reserved.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes (<http://creativecommons.org/licenses/by-nc/4.0/>).



Figure 1. Conventional X-rays showing erosions (white arrows) over right wrist, 4th proximal interphalangeal, and 3rd metacarpophalangeal joints with joint space narrowing over wrists.

resonance imaging. The patient was diagnosed with NMOSD associated with anti-AQP4-positive optic neuritis and an autoimmune disease RA.^{4,5}

She had refractory RA activity under MTX 15 mg/week, prednisolone 5 mg/day, and sulfasalazine (SAZ) 2 g/day. Conventional X-rays demonstrated erosions over right wrist, proximal interphalangeal and metacarpophalangeal joint areas with joint space narrowing over wrists (Figure 1). The RTX therapy with a regimen of 1 g × two fortnightly infusions every six months was initiated from 2016, with a total of 20 infusions until the end of 2020. There were completely depleted circulating CD19-positive B cells (0/μL) after two infusions, normalized ESR/CRP levels with a reduced DAS28 to 2.99 after four infusions, followed by negative anti-AQP4 (1 U/mL, normal reference below 3 U/mL), anti-CCP (below 7 U/mL), and RF (below 20 IU/mL) levels. She had an improved ophthalmological vision to decimal visual acuity 0.06/0.04 (right/left) in 2019. After 20 infusions, the DAS28 further improved to 2.10 and MTX doses were tapered to 7.5 mg/week without using SAZ. Except for initial infusion-related reactions, no adverse effects were observed during RTX therapeutic period. A written informed consent was

obtained from the patient for all diagnostic and therapeutic procedures.

Despite an extending spectrum of rheumatology disorders associated with NMO, including antiphospholipid syndrome, primary Sjögren's syndrome, SLE, and systemic sclerosis, there is no inclusion of RA.^{5,6} An investigation of NMOSD identified juvenile RA as a coexistent autoimmune disorder,⁷ and a survey on optic neuritis showed a poor visual outcome associated with RA.⁸ Interestingly, a case with overlap syndrome involving RA and polymyositis was diagnosed to have NMOSD with the presence of anti-AQP4-positive longitudinal myelopathy.⁹ Notably, in Han Chinese, genetic polymorphisms have been demonstrated to be shared among RA, NMOSD, and AQP4-seropositivity.¹⁰

In conclusion, in this report, we present an anti-AQP4-positive RA case with optic neuritis, which may raise a potential to incorporate RA into the autoimmunity spectrum in NMOSD.

Acknowledgments

The authors are indebted to physicians and nurses at our hospital involving in managing the reported patient. The Institutional Review Board approved this report (No. B-ER-105-108).

Declaration of conflicting interests

The authors declared no conflicts of interest with respect to the authorship and/or publication of this article.

Funding

The authors received no financial support for the research and/or authorship of this article.

REFERENCES

1. Wingerchuk DM, Lennon VA, Lucchinetti CF, Pittock SJ, Weinshenker BG. The spectrum of neuromyelitis optica. *Lancet Neurol* 2007;6:805-15.
2. Bennett JL, O'Connor KC, Bar-Or A, Zamvil SS, Hemmer B, Tedder TF, et al. B lymphocytes in neuromyelitis optica. *Neurol Neuroimmunol Neuroinflamm* 2015;2:e104.
3. Kim SH, Hyun JW, Kim HJ. Individualized B cell-targeting therapy for neuromyelitis optica spectrum disorder. *Neurochem Int* 2019;130:104347.
4. Wingerchuk DM, Banwell B, Bennett JL, Cabre P, Carroll W, Chitnis T, et al. International consensus diagnostic criteria for neuromyelitis optica spectrum disorders. *Neurology* 2015;85:177-89.
5. Freitas E, Guimarães J. Neuromyelitis optica spectrum disorders associated with other autoimmune diseases. *Rheumatol Int* 2015;35:243-53.
6. Lee HJ, Chang SH, Kang EH, Lee YJ, Song YW, Ha YJ. A case of primary Sjögren's syndrome presenting as mass-like encephalitis, with progression to neuromyelitis optica spectrum disorder. *Arch Rheumatol* 2017;32:353-7.
7. Pereira WLCJ, Reiche EMV, Kallaur AP, Oliveira SR, Simão ANC, Lozovoy MAB, et al. Frequency of autoimmune disorders and autoantibodies in patients with neuromyelitis optica. *Acta Neuropsychiatr* 2017;29:170-8.
8. Wang JC, Tow S, Aung T, Lim SA, Cullen JF. The presentation, aetiology, management and outcome of optic neuritis in an Asian population. *Clin Exp Ophthalmol* 2001;29:312-5.
9. Mizuno Y, Shinoda K, Watanabe M, Ogata H, Isobe N, Matsushita T, et al. Intractable axonal neuropathy with multifocal peripheral nerve swelling in neuromyelitis optica spectrum disorders: A case report. *Mult Scler Relat Disord* 2019;35:16-8.
10. Xie JL, Liu J, Lian ZY, Chen HX, Shi ZY, Zhang Q, et al. Association of GTF2IRD1-GTF2I polymorphisms with neuromyelitis optica spectrum disorders in Han Chinese patients. *Neural Regen Res* 2019;14:346-53.