

## Solitary syndesmophyte in odontoid process of a patient with ankylosing spondylitis

Hong Ki Min<sup>1</sup>, Se Hee Kim<sup>1</sup>, Hae-rim Kim<sup>2</sup>, Sang-heon Lee<sup>2</sup>

<sup>1</sup>Division of Rheumatology, Department of Internal Medicine, Konkuk University Medical Center, Seoul, Republic of Korea

<sup>2</sup>Division of Rheumatology, Department of Internal Medicine, Research Institute of Medical Science, Konkuk University School of Medicine, Seoul, Republic of Korea

A 22-year-old male patient was admitted to the rheumatology clinic due to neck and back pain with morning stiffness for 12 months. He suffered from right knee and heel pain six months ago. He showed increased levels of high-sensitivity C-reactive protein (2.20 mg/dL, reference range: 0.01-0.3 mg/dL) and erythrocyte sedimentation rate (67 mm/h, reference range: 0-15 mm/h). To evaluate neck and back pain, the physician performed computed tomography (CT) of the whole spine. The CT finding was absence for herniated nucleus pulposus nor neural foramen stenosis. All vertebral corners of whole spine were clear (Figure 1a), except for bony spur at odontoid process of C2 (Figure 1b, white arrow).

Upon suspicious of ankylosing spondylitis (AS), CT scan for sacroiliac joints and sacroiliac joint, and human leukocyte antigen (HLA)-B27 was tested. The CT scan showed bilateral Grade 2 sacroiliitis which was compatible with AS (Figure 1c), and positive result for HLA-B27. The

most widely used method for evaluating spinal structural damage, the modified Stoke Ankylosing Spondylitis Spinal Score only includes anterior aspect of C2 lower border to T1 upper border and T12 lower border to sacrum upper border,<sup>1</sup> and novel method using CT of whole spine (CT syndesmophyte score) includes lower border of C2 to upper border of sacrum.<sup>2</sup> These methods do not count the syndesmophyte on odontoid process of C2. The patient showed a rare case of AS which only showed syndesmophyte on odontoid process of C2.

Risks of osteoporosis and consequent vertebral fracture are increased in AS patients<sup>3</sup> and rarely fracture on odontoid process is reported.<sup>4</sup> Also, the low bone mineral density is associated with syndesmophyte progression in patients with AS.<sup>5</sup> Therefore, physicians should be aware of possibility of fracture or syndesmophyte progression on odontoid process when pain on upper cervical area occurs in AS patients.

**Received:** September 01, 2022 **Accepted:** September 29, 2022 **Published online:** November 04, 2022

**Correspondence:** Hong Ki Min, MD, PhD. Division of Rheumatology, Department of Internal Medicine, Konkuk University Medical Center, Seoul, 120-1, Neungdong-ro, Gwangjin-gu, Seoul, Republic of Korea.

e-mail: alsghdrl1921@naver.com

### Citation:

Min HK, Kim SH, Kim HR, Lee SH. Solitary syndesmophyte in odontoid process of a patient with ankylosing spondylitis. Arch Rheumatol 2023;38(x):i-ii.

©2023 Turkish League Against Rheumatism. All rights reserved.

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes (<http://creativecommons.org/licenses/by-nc/4.0/>).



**Figure 1.** Computed tomography images of (a) sacroiliac joint, (b) whole spine, and (c) syndesmophyte on odontoid process of C2 (white arrow).

**Ethics Committee Approval:** The present study was approved by the Institutional Review Board of Konkuk University Medical Center (KUMC 2022-07-002). The study was conducted in accordance with the principles of the Declaration of Helsinki.

**Patient Consent for Publication:** A written informed consent was obtained from the patient.

**Data Sharing Statement:** The data that support the findings of this study are available from the corresponding author upon reasonable request.

**Author Contributions:** Preparation and writing of the manuscript; data collection, analysis and interpretation; critical literature review; approval of the final version of the manuscript: H.K.M.; Data collection: S.H.K., H.R.K., S.H.Lee.

**Conflict of Interest:** The authors declared no conflicts of interest with respect to the authorship and/or publication of this article.

**Funding:** The authors received no financial support for the research and/or authorship of this article.

## REFERENCES

1. Creemers MC, Franssen MJ, van't Hof MA, Gribnau FW, van de Putte LB, van Riel PL. Assessment of outcome in ankylosing spondylitis: An extended radiographic scoring system. *Ann Rheum Dis* 2005;64:127-9.
2. de Bruin F, de Koning A, van den Berg R, Baraliakos X, Braun J, Ramiro S, et al. Development of the CT Syndesmophyte Score (CTSS) in patients with ankylosing spondylitis: Data from the SIAS cohort. *Ann Rheum Dis* 2018;77:371-7.
3. Magrey M, Khan MA. Osteoporosis in ankylosing spondylitis. *Curr Rheumatol Rep* 2010;12:332-6.
4. Eghbal K, Kamran H, Salimi A, Mamaghani HJ, Mirghaderi SP, Salimi M. Odontoid fracture complicating ankylosing spondylitis presenting with cervical canal stenosis and quadriplegia: A case report with 5-year follow-up and review of the literature. *Int J Surg Case Rep* 2022;94:107067.
5. Tan S, Bagheri H, Lee D, Shafiei A, Keaveny TM, Yao L, et al. Vertebral bone mineral density, vertebral strength, and syndesmophyte growth in ankylosing spondylitis: The importance of bridging. *Arthritis Rheumatol* 2022;74:1352-62.